



Role of Green Manuring in Improving Soil Fertility: A Sustainable Agricultural Approach

Maria Rodriguez ^{1*}, Emily Carter ², Jean-Pierre Dupont ³

¹ Lead Scientist, Agro-Fermentation Research Unit, National Agrarian University La Molina, Lima, Peru

² Assistant Professor, Department of Food Fermentation and Crop Sciences, University of Melbourne, Melbourne, Australia

³ Director, Institute of Agricultural and Food Fermentation, INRAE, Paris, France

* Corresponding Author: **Maria Rodriguez**

Article Info

Volume: 01

Issue: 01

January-February 2025

Received: 22-01-2025

Accepted: 19-02-2025

Page No: 14-17

Abstract

Green manuring represents a sustainable agricultural practice that significantly enhances soil fertility through the incorporation of living plant material into soil systems. This comprehensive review examines the role of green manuring in improving various aspects of soil fertility including nutrient availability, organic matter content, soil structure, and biological activity. The study analyzes different types of green manure crops, their mechanisms of action, and their impact on crop productivity and soil health. Findings indicate that green manuring practices can increase soil organic carbon by 15-30%, improve nitrogen availability by 40-80 kg/ha, and enhance soil physical properties substantially. However, successful implementation requires careful selection of appropriate green manure species, optimal timing of incorporation, and integration with existing cropping systems. The research concludes that green manuring is a cost-effective and environmentally sustainable approach to maintaining and improving soil fertility in agricultural systems.

Keywords: Green manuring, soil fertility, organic matter, nitrogen fixation, sustainable agriculture, soil health, cover crops

1. Introduction

Soil fertility decline represents one of the most significant challenges facing modern agriculture, with intensive farming practices leading to degradation of soil physical, chemical, and biological properties¹. The excessive reliance on synthetic fertilizers has created economic and environmental concerns, necessitating the adoption of sustainable soil management practices². Green manuring emerges as a promising solution that addresses soil fertility challenges while promoting environmental sustainability³. Green manuring involves the cultivation of specific crops that are subsequently incorporated into the soil to improve fertility and soil health⁴. This ancient practice has gained renewed attention due to its multiple benefits including nitrogen fixation, organic matter addition, soil structure improvement, and pest management⁵. The integration of green manuring with modern agricultural systems offers opportunities to reduce external input dependence while maintaining or increasing crop productivity⁶. The practice encompasses various approaches including dedicated green manure crops, cover crops, and intercropping systems that contribute organic matter and nutrients to soil systems⁷. Understanding the mechanisms and benefits of green manuring is essential for developing sustainable agricultural practices that can meet food security needs while preserving soil resources for future generations⁸.

2. Types and Classification of Green Manure Crops

2.1 Leguminous Green Manure Crops

Leguminous crops constitute the most important category of green manure plants due to their nitrogen-fixing capabilities through symbiotic relationships with rhizobia bacteria⁹. Common leguminous green manure crops include dhaincha (*Sesbania aculeata*),

sun hemp (*Crotalaria juncea*), cowpea (*Vigna unguiculata*), and cluster bean (*Cyamopsis tetragonoloba*)¹⁰. These crops can fix atmospheric nitrogen ranging from 50-300 kg/ha depending on species, growing conditions, and management practices¹¹. *Sesbania* species are particularly valuable due to their rapid growth, high biomass production, and ability to thrive in waterlogged conditions¹². Sun hemp demonstrates excellent performance in tropical and subtropical regions, producing substantial biomass while fixing significant amounts of nitrogen¹³. The selection of appropriate leguminous species depends on local climate conditions, soil types, and cropping system requirements¹⁴.

2.2 Non-leguminous Green Manure Crops

Non-leguminous green manure crops contribute primarily through biomass addition and nutrient recycling rather than nitrogen fixation¹⁵. Important species include mustard (*Brassica* species), oats (*Avena sativa*), rye (*Secale cereale*), and various grasses¹⁶. These crops excel in scavenging nutrients from deeper soil layers, preventing nutrient leaching, and providing diverse organic matter inputs¹⁷.

Brassicaceae family crops possess unique characteristics including deep root systems that break soil compaction and release glucosinolates that exhibit biofumigation properties¹⁸. Cereal green manures contribute high carbon content organic matter that improves soil structure and provides energy sources for soil microorganisms¹⁹. The integration of leguminous and non-leguminous species creates balanced green manure systems with complementary benefits²⁰.

3. Mechanisms of Soil Fertility Improvement

3.1 Biological Nitrogen Fixation

The primary mechanism through which leguminous green manures improve soil fertility is biological nitrogen fixation²¹. Rhizobia bacteria form symbiotic relationships with legume roots, converting atmospheric nitrogen into plant-available forms²². This process reduces dependence on synthetic nitrogen fertilizers while providing sustained nitrogen release through decomposition²³.

The efficiency of nitrogen fixation depends on various factors including rhizobia strain effectiveness, soil pH, moisture conditions, and nutrient availability²⁴. Optimal conditions can result in nitrogen fixation rates of 100-300 kg/ha annually, equivalent to substantial fertilizer inputs²⁵. The fixed nitrogen becomes available gradually as green manure decomposes, providing sustained nutrient supply throughout the growing season²⁶.

3.2 Organic Matter Addition and Decomposition

Green manuring significantly increases soil organic matter content through the incorporation of fresh plant material²⁷. The decomposition of green manure releases nutrients gradually while contributing to soil humus formation²⁸. Studies indicate that regular green manuring can increase soil organic carbon by 0.2-0.5% annually, substantially improving soil fertility²⁹.

The decomposition process involves complex microbial activities that transform plant materials into stable organic compounds³⁰. Different plant tissues decompose at varying rates, with leaves and soft stems decomposing rapidly while woody materials contribute to long-term organic matter pools³¹. The carbon-to-nitrogen ratio of green manure materials influences decomposition rates and nutrient release

patterns³².

3.3 Nutrient Cycling and Availability

Green manure crops enhance nutrient cycling by accessing nutrients from various soil layers and concentrating them in above-ground biomass³³. Deep-rooted species can retrieve nutrients from subsoil layers that are otherwise unavailable to shallow-rooted crops³⁴. Upon incorporation, these nutrients become available in the surface soil where most crop roots are concentrated³⁵.

Phosphorus availability often increases following green manuring due to organic acid production during decomposition and enhanced microbial activity³⁶. Potassium and micronutrients are also recycled effectively through green manure systems³⁷. The gradual release of nutrients from decomposing organic matter provides sustained nutrient availability that matches crop uptake patterns³⁸.

4. Impact on Soil Physical Properties

4.1 Soil Structure and Aggregation

Green manuring significantly improves soil structure through increased organic matter content and enhanced biological activity³⁹. Soil aggregation increases substantially following green manure incorporation, improving soil porosity and water infiltration⁴⁰. Studies demonstrate that green manuring can increase soil aggregation by 25-40% compared to conventional practices⁴¹.

Root systems of green manure crops create channels and macropores that persist after decomposition, improving soil aeration and water movement⁴². The production of binding agents during decomposition helps stabilize soil aggregates against erosion and compaction⁴³. Enhanced soil structure benefits include improved seedbed preparation, better root penetration, and reduced soil crusting⁴⁴.

4.2 Water Retention and Infiltration

Improved organic matter content from green manuring enhances soil water retention capacity⁴⁵. Organic matter can hold up to 20 times its weight in water, significantly increasing soil moisture availability for crops⁴⁶. This increased water holding capacity is particularly beneficial in rainfed agriculture and water-scarce regions⁴⁷.

Water infiltration rates improve substantially following green manuring due to enhanced soil structure and porosity⁴⁸. Reduced surface runoff and increased water storage contribute to improved drought resilience and reduced irrigation requirements⁴⁹. The combination of improved infiltration and retention creates favorable water regimes for crop growth⁵⁰.

5. Enhancement of Soil Biological Activity

5.1 Microbial Population and Diversity

Green manuring significantly enhances soil microbial populations and diversity through increased organic matter inputs⁵¹. Microbial biomass typically increases by 30-60% following green manure incorporation⁵². The diverse organic compounds in green manure support various microbial communities with different functional roles⁵³.

Enhanced microbial activity contributes to nutrient cycling, organic matter decomposition, and soil structure formation⁵⁴. Beneficial microorganisms including mycorrhizal fungi, nitrogen-fixing bacteria, and plant growth-promoting rhizobacteria increase substantially in green manured soils⁵⁵.

This enhanced biological activity creates a more resilient and sustainable soil ecosystem⁵⁶.

5.2 Enzyme Activity and Biochemical Processes

Soil enzyme activities increase significantly following green manuring, indicating enhanced biochemical processes⁵⁷. Important enzymes including urease, phosphatase, and dehydrogenase show substantial increases in green manured soils⁵⁸. These enzymes catalyze essential nutrient transformation processes that improve nutrient availability⁵⁹. The enhanced enzymatic activity reflects increased microbial metabolism and organic matter turnover⁶⁰. This increased biological activity contributes to soil health and resilience against stress factors⁶¹. The biochemical improvements from green manuring have long-lasting effects that benefit subsequent crops⁶².

6. Effects on Crop Productivity and Quality

6.1 Yield Improvements

Numerous studies demonstrate significant yield improvements following green manuring across various crops and agro-climatic conditions⁶³. Rice yields typically increase by 15-25% when preceded by green manuring with leguminous crops⁶⁴. Wheat and maize also show substantial yield responses to green manuring, particularly in low-fertility soils⁶⁵.

The yield improvements result from multiple factors including improved nutrient availability, enhanced soil physical properties, and better water relations⁶⁶. Long-term studies indicate that yield benefits from green manuring are sustained over multiple cropping seasons⁶⁷. The cumulative effects of green manuring become more pronounced with continued practice⁶⁸.

6.2 Crop Quality Enhancement

Green manuring improves crop quality parameters including protein content, nutritional value, and processing characteristics⁶⁹. The gradual nutrient release from green manure supports balanced crop nutrition that enhances quality attributes⁷⁰. Improved soil biological activity also contributes to better nutrient uptake and crop health⁷¹.

Studies indicate that crops grown following green manuring show improved resistance to pests and diseases⁷². The enhanced soil health and plant nutrition contribute to stronger plant defense mechanisms⁷³. Quality improvements from green manuring can result in premium prices and better market acceptance⁷⁴.

7. Economic and Environmental Benefits

7.1 Economic Advantages

Green manuring provides significant economic benefits through reduced fertilizer costs and improved crop yields⁷⁵. Studies indicate that green manuring can reduce nitrogen fertilizer requirements by 25-50% while maintaining or increasing yields⁷⁶. The cost of green manuring is typically lower than equivalent fertilizer inputs⁷⁷.

Long-term economic benefits include improved soil health, reduced input costs, and sustained productivity⁷⁸. The practice also provides additional income opportunities through seed production and biomass utilization⁷⁹. Risk reduction through improved soil fertility and drought resilience adds to the economic value⁸⁰.

7.2 Environmental Benefits

Environmental benefits of green manuring include reduced greenhouse gas emissions, improved carbon sequestration, and decreased environmental pollution⁸¹. Carbon sequestration in green manured soils can range from 0.5-2.0 tonnes CO₂ equivalent per hectare annually⁸². Reduced synthetic fertilizer use decreases energy consumption and associated emissions⁸³.

Water quality improvements result from reduced fertilizer runoff and improved soil stability⁸⁴. Biodiversity enhancement occurs through habitat provision and reduced pesticide use⁸⁵. The practice contributes to sustainable agricultural systems that maintain ecosystem services⁸⁶.

8. Challenges and Limitations

8.1 Management Challenges

Successful green manuring requires careful planning and management to optimize benefits⁸⁷. Timing of incorporation affects decomposition rates and nutrient availability⁸⁸. Weather conditions during growth and incorporation significantly influence effectiveness⁸⁹.

Water requirements for green manure crops may compete with cash crops in water-scarce regions⁹⁰. Labor requirements for establishment and incorporation can be substantial⁹¹. Integration with existing cropping systems requires modifications to traditional practices⁹².

8.2 Limitations and Constraints

Initial establishment costs and opportunity costs of land use present barriers to adoption⁹³. Slow nutrient release may not meet immediate crop requirements⁹⁴. Variability in performance across different soil types and climatic conditions creates uncertainty⁹⁵.

Pest and disease issues associated with some green manure crops require careful management⁹⁶. Market limitations for green manure seeds and lack of technical knowledge hinder widespread adoption⁹⁷. Policy support and extension services are often inadequate for promoting green manuring practices⁹⁸.

9. Future Prospects and Recommendations

9.1 Research Priorities

Future research should focus on developing improved green manure varieties with enhanced nitrogen fixation and biomass production⁹⁹. Integration with precision agriculture technologies can optimize green manuring practices¹⁰⁰. Long-term studies are needed to quantify cumulative benefits and sustainability¹⁰¹.

Climate change impacts on green manuring effectiveness require investigation¹⁰². Economic analyses considering all costs and benefits will support policy development¹⁰³. Participatory research approaches can improve farmer adoption and adaptation¹⁰⁴.

9.2 Policy and Extension Recommendations

Government policies should provide incentives for green manuring adoption including subsidies and technical support¹⁰⁵. Extension programs must educate farmers about benefits and best practices¹⁰⁶. Research and development investments should focus on location-specific recommendations¹⁰⁷.

Market development for green manure seeds and equipment will support widespread adoption¹⁰⁸. Integration with existing

agricultural programs can mainstream green manuring practices¹⁰⁹. International cooperation can facilitate technology transfer and knowledge sharing¹¹⁰.

10. Conclusion

Green manuring represents a scientifically proven and economically viable approach to improving soil fertility in agricultural systems. The practice offers multiple benefits including enhanced nutrient availability, improved soil physical and biological properties, and increased crop productivity. The mechanisms of action are well understood, involving biological nitrogen fixation, organic matter addition, and enhanced nutrient cycling.

The evidence clearly demonstrates that green manuring can significantly improve soil fertility while providing economic and environmental benefits. Yield improvements of 15-25% are commonly achieved, with additional benefits including improved crop quality and reduced input costs. The practice contributes to sustainable agriculture by reducing dependence on synthetic fertilizers while maintaining productivity.

However, successful implementation requires addressing management challenges and providing adequate support systems. Careful selection of appropriate species, optimal timing of operations, and integration with existing farming systems are essential for success. Policy support, extension services, and market development are needed to promote widespread adoption.

The future of green manuring lies in developing improved varieties, integrating with modern technologies, and creating supportive policy environments. Long-term research is needed to fully understand the cumulative benefits and optimize practices for different agro-ecological conditions. With appropriate support and management, green manuring can play a crucial role in achieving sustainable and productive agricultural systems.

Green manuring offers a path toward sustainable intensification of agriculture that meets food security needs while preserving soil resources and environmental quality. The practice aligns with global sustainability goals and climate change mitigation efforts. Continued research, policy support, and farmer education will be essential for realizing the full potential of green manuring in improving soil fertility and agricultural sustainability.

11. References

1. Lal R. Soil organic matter content and crop yields in tropical and temperate regions. *Soil Use Manag.* 2020;36(4):663–78.
2. Singh Y, Sidhu HS. Management of cereal crop residues for sustainable rice-wheat production system in the Indo-Gangetic plains of India. *Proc Indian Natl Sci Acad.* 2014;80(1):95–114.
3. Fageria NK. Green manuring in crop production. *J Plant Nutr.* 2007;30(5):691–719.
4. Kumar S, Dhar S. Effect of green manuring on soil properties and crop yield in rice-wheat cropping system. *J Indian Soc Soil Sci.* 2010;58(4):409–12.
5. Cherr CM, Scholberg JM, McSorley R. Green manure approaches to crop production: A synthesis. *Agron J.* 2006;98(2):302–19.
6. Palm CA, Myers RJK, Nandwa SM. Organic inputs for soil fertility management in tropical agroecosystems:

Application of an organic resource database. *Agric Ecosyst Environ.* 2001;83(1–2):27–42.

7. Snapp SS, Blackie MJ, Gilbert RA, Bezner-Kerr R, Kanyama-Phiri GY. Biodiversity can support a greener revolution in Africa. *Proc Natl Acad Sci U S A.* 2010;107(48):20840–5.
8. Tilman D, Balzer C, Hill J, Befort BL. Global food demand and the sustainable intensification of agriculture. *Proc Natl Acad Sci U S A.* 2011;108(50):20260–4.
9. Peoples MB, Herridge DF, Ladha JK. The contributions of nitrogen-fixing crop legumes to the productivity of agricultural systems. *Symbiosis.* 2009;48(1–3):1–17.
10. Becker M, Johnson DE. Improved water control and crop management effects on lowland rice productivity in West Africa. *Agron J.* 2001;93(6):1322–31.